

**APPLICATION  
FOR  
UNITED STATES LETTERS PATENT**

**TITLE: TREATMENT FOR DIABETES**

**INVENTOR(S): INDU PARIKH, ANNE LANE,  
RONALD V. NARDI and  
STEPHEN J. BRAND**

**CERTIFICATE OF EXPRESS MAILING**

"Express Mail" Label No.: EL79606266649  
Date of Deposit: Dec 20, 2001

I hereby certify under 37 C.F.R. 1.10 that this correspondence is being deposited with the United States Postal Service as "Express Mail Post Office to Addressee" with sufficient postage on the date indicated above and is addressed to the Commissioner of Patents and Trademarks, PO Box 2327 Arlington, VA 22202.

Robert Pattison  
(Signature)  
Robert Pattison

**TREATMENT FOR DIABETES****CROSS-REFERENCE TO RELATED APPLICATIONS**

5 Sub ~~Q~~ This application is a continuation-in-part of ~~USSN 09/127,028, filed July 30, 1998,~~  
which is a continuation of ~~USSN 07/992,255, filed December 14, 1992, which issued~~  
March 23, 1999, as USPN \_\_\_\_\_, which disclosures are incorporated herein by  
reference.

**INTRODUCTION****Field of Invention**

10  
15  
20  
25  
30  
35  
40  
45  
50  
55  
60  
65  
70  
75  
80  
85  
90  
95  
100  
105  
110  
115  
120  
125  
130  
135  
140  
145  
150  
155  
160  
165  
170  
175  
180  
185  
190  
195  
200  
205  
210  
215  
220  
225  
230  
235  
240  
245  
250  
255  
260  
265  
270  
275  
280  
285  
290  
295  
300  
305  
310  
315  
320  
325  
330  
335  
340  
345  
350  
355  
360  
365  
370  
375  
380  
385  
390  
395  
400  
405  
410  
415  
420  
425  
430  
435  
440  
445  
450  
455  
460  
465  
470  
475  
480  
485  
490  
495  
500  
505  
510  
515  
520  
525  
530  
535  
540  
545  
550  
555  
560  
565  
570  
575  
580  
585  
590  
595  
600  
605  
610  
615  
620  
625  
630  
635  
640  
645  
650  
655  
660  
665  
670  
675  
680  
685  
690  
695  
700  
705  
710  
715  
720  
725  
730  
735  
740  
745  
750  
755  
760  
765  
770  
775  
780  
785  
790  
795  
800  
805  
810  
815  
820  
825  
830  
835  
840  
845  
850  
855  
860  
865  
870  
875  
880  
885  
890  
895  
900  
905  
910  
915  
920  
925  
930  
935  
940  
945  
950  
955  
960  
965  
970  
975  
980  
985  
990  
995  
1000  
1005  
1010  
1015  
1020  
1025  
1030  
1035  
1040  
1045  
1050  
1055  
1060  
1065  
1070  
1075  
1080  
1085  
1090  
1095  
1100  
1105  
1110  
1115  
1120  
1125  
1130  
1135  
1140  
1145  
1150  
1155  
1160  
1165  
1170  
1175  
1180  
1185  
1190  
1195  
1200  
1205  
1210  
1215  
1220  
1225  
1230  
1235  
1240  
1245  
1250  
1255  
1260  
1265  
1270  
1275  
1280  
1285  
1290  
1295  
1300  
1305  
1310  
1315  
1320  
1325  
1330  
1335  
1340  
1345  
1350  
1355  
1360  
1365  
1370  
1375  
1380  
1385  
1390  
1395  
1400  
1405  
1410  
1415  
1420  
1425  
1430  
1435  
1440  
1445  
1450  
1455  
1460  
1465  
1470  
1475  
1480  
1485  
1490  
1495  
1500  
1505  
1510  
1515  
1520  
1525  
1530  
1535  
1540  
1545  
1550  
1555  
1560  
1565  
1570  
1575  
1580  
1585  
1590  
1595  
1600  
1605  
1610  
1615  
1620  
1625  
1630  
1635  
1640  
1645  
1650  
1655  
1660  
1665  
1670  
1675  
1680  
1685  
1690  
1695  
1700  
1705  
1710  
1715  
1720  
1725  
1730  
1735  
1740  
1745  
1750  
1755  
1760  
1765  
1770  
1775  
1780  
1785  
1790  
1795  
1800  
1805  
1810  
1815  
1820  
1825  
1830  
1835  
1840  
1845  
1850  
1855  
1860  
1865  
1870  
1875  
1880  
1885  
1890  
1895  
1900  
1905  
1910  
1915  
1920  
1925  
1930  
1935  
1940  
1945  
1950  
1955  
1960  
1965  
1970  
1975  
1980  
1985  
1990  
1995  
2000  
2005  
2010  
2015  
2020  
2025  
2030  
2035  
2040  
2045  
2050  
2055  
2060  
2065  
2070  
2075  
2080  
2085  
2090  
2095  
2100  
2105  
2110  
2115  
2120  
2125  
2130  
2135  
2140  
2145  
2150  
2155  
2160  
2165  
2170  
2175  
2180  
2185  
2190  
2195  
2200  
2205  
2210  
2215  
2220  
2225  
2230  
2235  
2240  
2245  
2250  
2255  
2260  
2265  
2270  
2275  
2280  
2285  
2290  
2295  
2300  
2305  
2310  
2315  
2320  
2325  
2330  
2335  
2340  
2345  
2350  
2355  
2360  
2365  
2370  
2375  
2380  
2385  
2390  
2395  
2400  
2405  
2410  
2415  
2420  
2425  
2430  
2435  
2440  
2445  
2450  
2455  
2460  
2465  
2470  
2475  
2480  
2485  
2490  
2495  
2500  
2505  
2510  
2515  
2520  
2525  
2530  
2535  
2540  
2545  
2550  
2555  
2560  
2565  
2570  
2575  
2580  
2585  
2590  
2595  
2600  
2605  
2610  
2615  
2620  
2625  
2630  
2635  
2640  
2645  
2650  
2655  
2660  
2665  
2670  
2675  
2680  
2685  
2690  
2695  
2700  
2705  
2710  
2715  
2720  
2725  
2730  
2735  
2740  
2745  
2750  
2755  
2760  
2765  
2770  
2775  
2780  
2785  
2790  
2795  
2800  
2805  
2810  
2815  
2820  
2825  
2830  
2835  
2840  
2845  
2850  
2855  
2860  
2865  
2870  
2875  
2880  
2885  
2890  
2895  
2900  
2905  
2910  
2915  
2920  
2925  
2930  
2935  
2940  
2945  
2950  
2955  
2960  
2965  
2970  
2975  
2980  
2985  
2990  
2995  
3000  
3005  
3010  
3015  
3020  
3025  
3030  
3035  
3040  
3045  
3050  
3055  
3060  
3065  
3070  
3075  
3080  
3085  
3090  
3095  
3100  
3105  
3110  
3115  
3120  
3125  
3130  
3135  
3140  
3145  
3150  
3155  
3160  
3165  
3170  
3175  
3180  
3185  
3190  
3195  
3200  
3205  
3210  
3215  
3220  
3225  
3230  
3235  
3240  
3245  
3250  
3255  
3260  
3265  
3270  
3275  
3280  
3285  
3290  
3295  
3300  
3305  
3310  
3315  
3320  
3325  
3330  
3335  
3340  
3345  
3350  
3355  
3360  
3365  
3370  
3375  
3380  
3385  
3390  
3395  
3400  
3405  
3410  
3415  
3420  
3425  
3430  
3435  
3440  
3445  
3450  
3455  
3460  
3465  
3470  
3475  
3480  
3485  
3490  
3495  
3500  
3505  
3510  
3515  
3520  
3525  
3530  
3535  
3540  
3545  
3550  
3555  
3560  
3565  
3570  
3575  
3580  
3585  
3590  
3595  
3600  
3605  
3610  
3615  
3620  
3625  
3630  
3635  
3640  
3645  
3650  
3655  
3660  
3665  
3670  
3675  
3680  
3685  
3690  
3695  
3700  
3705  
3710  
3715  
3720  
3725  
3730  
3735  
3740  
3745  
3750  
3755  
3760  
3765  
3770  
3775  
3780  
3785  
3790  
3795  
3800  
3805  
3810  
3815  
3820  
3825  
3830  
3835  
3840  
3845  
3850  
3855  
3860  
3865  
3870  
3875  
3880  
3885  
3890  
3895  
3900  
3905  
3910  
3915  
3920  
3925  
3930  
3935  
3940  
3945  
3950  
3955  
3960  
3965  
3970  
3975  
3980  
3985  
3990  
3995  
4000  
4005  
4010  
4015  
4020  
4025  
4030  
4035  
4040  
4045  
4050  
4055  
4060  
4065  
4070  
4075  
4080  
4085  
4090  
4095  
4100  
4105  
4110  
4115  
4120  
4125  
4130  
4135  
4140  
4145  
4150  
4155  
4160  
4165  
4170  
4175  
4180  
4185  
4190  
4195  
4200  
4205  
4210  
4215  
4220  
4225  
4230  
4235  
4240  
4245  
4250  
4255  
4260  
4265  
4270  
4275  
4280  
4285  
4290  
4295  
4300  
4305  
4310  
4315  
4320  
4325  
4330  
4335  
4340  
4345  
4350  
4355  
4360  
4365  
4370  
4375  
4380  
4385  
4390  
4395  
4400  
4405  
4410  
4415  
4420  
4425  
4430  
4435  
4440  
4445  
4450  
4455  
4460  
4465  
4470  
4475  
4480  
4485  
4490  
4495  
4500  
4505  
4510  
4515  
4520  
4525  
4530  
4535  
4540  
4545  
4550  
4555  
4560  
4565  
4570  
4575  
4580  
4585  
4590  
4595  
4600  
4605  
4610  
4615  
4620  
4625  
4630  
4635  
4640  
4645  
4650  
4655  
4660  
4665  
4670  
4675  
4680  
4685  
4690  
4695  
4700  
4705  
4710  
4715  
4720  
4725  
4730  
4735  
4740  
4745  
4750  
4755  
4760  
4765  
4770  
4775  
4780  
4785  
4790  
4795  
4800  
4805  
4810  
4815  
4820  
4825  
4830  
4835  
4840  
4845  
4850  
4855  
4860  
4865  
4870  
4875  
4880  
4885  
4890  
4895  
4900  
4905  
4910  
4915  
4920  
4925  
4930  
4935  
4940  
4945  
4950  
4955  
4960  
4965  
4970  
4975  
4980  
4985  
4990  
4995  
5000  
5005  
5010  
5015  
5020  
5025  
5030  
5035  
5040  
5045  
5050  
5055  
5060  
5065  
5070  
5075  
5080  
5085  
5090  
5095  
5100  
5105  
5110  
5115  
5120  
5125  
5130  
5135  
5140  
5145  
5150  
5155  
5160  
5165  
5170  
5175  
5180  
5185  
5190  
5195  
5200  
5205  
5210  
5215  
5220  
5225  
5230  
5235  
5240  
5245  
5250  
5255  
5260  
5265  
5270  
5275  
5280  
5285  
5290  
5295  
5300  
5305  
5310  
5315  
5320  
5325  
5330  
5335  
5340  
5345  
5350  
5355  
5360  
5365  
5370  
5375  
5380  
5385  
5390  
5395  
5400  
5405  
5410  
5415  
5420  
5425  
5430  
5435  
5440  
5445  
5450  
5455  
5460  
5465  
5470  
5475  
5480  
5485  
5490  
5495  
5500  
5505  
5510  
5515  
5520  
5525  
5530  
5535  
5540  
5545  
5550  
5555  
5560  
5565  
5570  
5575  
5580  
5585  
5590  
5595  
5600  
5605  
5610  
5615  
5620  
5625  
5630  
5635  
5640  
5645  
5650  
5655  
5660  
5665  
5670  
5675  
5680  
5685  
5690  
5695  
5700  
5705  
5710  
5715  
5720  
5725  
5730  
5735  
5740  
5745  
5750  
5755  
5760  
5765  
5770  
5775  
5780  
5785  
5790  
5795  
5800  
5805  
5810  
5815  
5820  
5825  
5830  
5835  
5840  
5845  
5850  
5855  
5860  
5865  
5870  
5875  
5880  
5885  
5890  
5895  
5900  
5905  
5910  
5915  
5920  
5925  
5930  
5935  
5940  
5945  
5950  
5955  
5960  
5965  
5970  
5975  
5980  
5985  
5990  
5995  
6000  
6005  
6010  
6015  
6020  
6025  
6030  
6035  
6040  
6045  
6050  
6055  
6060  
6065  
6070  
6075  
6080  
6085  
6090  
6095  
6100  
6105  
6110  
6115  
6120  
6125  
6130  
6135  
6140  
6145  
6150  
6155  
6160  
6165  
6170  
6175  
6180  
6185  
6190  
6195  
6200  
6205  
6210  
6215  
6220  
6225  
6230  
6235  
6240  
6245  
6250  
6255  
6260  
6265  
6270  
6275  
6280  
6285  
6290  
6295  
6300  
6305  
6310  
6315  
6320  
6325  
6330  
6335  
6340  
6345  
6350  
6355  
6360  
6365  
6370  
6375  
6380  
6385  
6390  
6395  
6400  
6405  
6410  
6415  
6420  
6425  
6430  
6435  
6440  
6445  
6450  
6455  
6460  
6465  
6470  
6475  
6480  
6485  
6490  
6495  
6500  
6505  
6510  
6515  
6520  
6525  
6530  
6535  
6540  
6545  
6550  
6555  
6560  
6565  
6570  
6575  
6580  
6585  
6590  
6595  
6600  
6605  
6610  
6615  
6620  
6625  
6630  
6635  
6640  
6645  
6650  
6655  
6660  
6665  
6670  
6675  
6680  
6685  
6690  
6695  
6700  
6705  
6710  
6715  
6720  
6725  
6730  
6735  
6740  
6745  
6750  
6755  
6760  
6765  
6770  
6775  
6780  
6785  
6790  
6795  
6800  
6805  
6810  
6815  
6820  
6825  
6830  
6835  
6840  
6845  
6850  
6855  
6860  
6865  
6870  
6875  
6880  
6885  
6890  
6895  
6900  
6905  
6910  
6915  
6920  
6925  
6930  
6935  
6940  
6945  
6950  
6955  
6960  
6965  
6970  
6975  
6980  
6985  
6990  
6995  
7000  
7005  
7010  
7015  
7020  
7025  
7030  
7035  
7040  
7045  
7050  
7055  
7060  
7065  
7070  
7075  
7080  
7085  
7090  
7095  
7100  
7105  
7110  
7115  
7120  
7125  
7130  
7135  
7140  
7145  
7150  
7155  
7160  
7165  
7170  
7175  
7180  
7185  
7190  
7195  
7200  
7205  
7210  
7215  
7220  
7225  
7230  
7235  
7240  
7245  
7250  
7255  
7260  
7265  
7270  
7275  
7280  
7285  
7290  
7295  
7300  
7305  
7310  
7315  
7320  
7325  
7330  
7335  
7340  
7345  
7350  
7355  
7360  
7365  
7370  
7375  
7380  
7385  
7390  
7395  
7400  
7405  
7410  
7415  
7420  
7425  
7430  
7435  
7440  
7445  
7450  
7455  
7460  
7465  
7470  
7475  
7480  
7485  
7490  
7495  
7500  
7505  
7510  
7515  
7520  
7525  
7530  
7535  
7540  
7545  
7550  
7555  
7560  
7565  
7570  
7575  
7580  
7585  
7590  
7595  
7600  
7605  
7610  
7615  
7620  
7625  
7630  
7635  
7640  
7645  
7650  
7655  
7660  
7665  
7670  
7675  
7680  
7685  
7690  
7695  
7700  
7705  
7710  
7715  
7720  
7725  
7730  
7735  
7740  
7745  
7750  
7755  
7760  
7765  
7770  
7775  
7780  
7785  
7790  
7795  
7800  
7805  
7810  
7815  
7820  
7825  
7830  
7835  
7840  
7845  
7850  
7855  
7860  
7865  
7870  
7875  
7880  
7885  
7890  
7895  
7900  
7905  
7910  
7915  
7920  
7925  
7930  
7935  
7940  
7945  
7950  
7955  
7960  
7965  
7970  
7975  
7980  
7985  
7990  
7995  
8000  
8005  
8010  
8015  
8020  
8025  
8030  
8035  
8040  
8045  
8050  
8055  
8060  
8065  
8070  
8075  
8080  
8085  
8090  
8095  
8100  
8105  
8110  
8115  
8120  
8125  
8130  
8135  
8140  
8145  
8150  
8155  
8160  
8165  
8170  
8175  
8180  
8185  
8190  
8195  
8200  
8205  
8210  
8215  
8220  
8225  
8230  
8235  
8240  
8245  
8250  
8255  
8260  
8265  
8270  
8275  
8280  
8285  
8290  
8295  
8300  
8305  
8310  
8315  
8320  
8325  
8330  
8335  
8340  
8345  
8350  
8355  
8360  
8365  
8370  
8375  
8380  
8385  
8390  
8395  
8400  
8405  
8410  
8415  
8420  
8425  
8430  
8435  
8440  
8445  
8450  
8455  
8460  
8465  
8470  
8475  
8480  
8485  
8490  
8495  
8500  
8505  
8510  
8515  
8520  
8525  
8530  
8535  
8540  
8545  
8550  
8555  
8560  
8565  
8570  
8575  
8580  
8585  
8590  
8595  
8600  
8605  
8610  
8615  
8620  
8625  
8630  
8635  
8640  
8645  
8650  
8655  
8660  
8665  
8670  
8675  
8680  
8685  
8690  
8695  
8700  
8705  
8710  
8715  
8720  
8725  
8730  
8735  
8740  
8745  
8750  
8755  
8760  
8765  
8770  
8775  
8780  
8785  
8790  
8795  
8800  
8805  
8810  
8815  
8820  
8825  
8830  
8835  
8840  
8845  
8850  
8855  
8860  
8865  
8870  
8875  
8880  
8885  
8890  
8895  
8900  
8905  
8910  
8915  
8920  
8925  
8930  
8935  
8940  
8945  
8950  
8955  
8960  
8965  
8970  
8975  
8980  
8985  
8990  
8995  
9000  
9005  
9010  
9015  
9020  
9025  
9030  
9035  
9040  
9045  
9050  
9055  
9060  
9065  
9070  
9075  
9080  
9085  
9090  
9095  
9100  
9105  
9110  
9115  
9120  
9125  
9130  
9135  
9140  
9145  
9150  
9155  
9160  
9165  
9170  
9175  
9180  
9185  
9190  
9195  
9200  
9205  
9210  
9215  
9220  
9225  
9230  
9235  
9240  
9245  
9250  
9255  
9260  
9265  
9270  
9275  
9280  
9285  
9290  
9295  
9300  
9305  
9310  
9315  
9320  
9325  
9330  
9335  
9340  
9345  
9350  
9355  
9360  
9365  
9370  
9375  
9380  
9385  
9390  
9395  
9400  
9405  
9410  
9415  
9420  
9425  
9430  
9435  
9440  
9445  
9450  
9455  
9460  
9465  
9470  
9475  
9480  
9485  
9490  
9495  
9500  
9505  
9510  
9515  
9520  
9525  
9530  
9535  
9540  
9545  
9550  
9555  
9560  
9565  
9570  
9575  
9580  
9585  
9590  
9595  
9600  
9605  
9610  
9615  
9620  
9625  
9630  
9635  
9640  
9645  
9650  
9655  
9660  
9665  
9670  
9675  
9680  
9685  
9690  
9695  
9700  
9705  
9710  
9715  
9720  
9725  
9730  
9735  
9740  
9745  
9750  
9755  
9760  
9765  
9770  
9775  
9780  
9785  
9790  
9795  
9800  
9805  
9810  
9815  
9820  
9825  
9830  
9835  
9840  
9845  
9850  
9855  
9860  
9865  
9870  
9875  
9880  
9885  
9890  
9895  
9900  
9905  
9910  
9915  
9920  
9925  
9930  
9935  
9940  
9945  
9950  
9955  
9960  
9965  
9970  
9975  
9980  
9985  
9990  
9995  
10000  
10005  
10010  
10015  
10020  
10025  
10030  
10035  
10040  
10045  
10050  
10055  
10060  
10065  
10070  
10075  
10080  
10085  
10090  
10095  
10100  
10105  
1011

metabolic control. Initial drug therapy includes sulfonylureas that stimulate  $\beta$ -cell insulin secretion, but also can include biguanides,  $\alpha$ -glucosidase inhibitors, thiazolidenediones and combination therapy. It is noteworthy however that the progressive nature of the disease mechanisms operating in type 2 diabetes are difficult to control. Over 50% of all drug-treated diabetics demonstrate poor glycemic control within six years, irrespective of the drug administered. Insulin therapy is regarded by many as the last resort in the treatment of Type 2 diabetes, and there is patient resistance to the use of insulin.

Pancreatic islets develop from endodermal stem cells that lie in the fetal ductular pancreatic endothelium, which also contains pluripotent stem cells that develop into the exocrine pancreas. Teitelman and Lee, *Developmental Biology*, 121:454-466 (1987); Pictet and Rutter, *Development of the embryonic endocrine pancreas, in Endocrinology, Handbook of Physiology*, ed. R.O. Greep and E.B. Astwood (1972), American Physiological Society: Washington, D.C., p.25-66. Islet development proceeds through discrete developmental stages during fetal gestation which are punctuated by dramatic transitions. The initial period is a protodifferentiated state which is characterized by the commitment of the pluripotent stem cells to the islet cell lineage, as manifested by the expression of insulin and glucagon by the protodifferentiated cells. These protodifferentiated cells comprise a population of committed islet precursor cells which express only low levels of islet specific gene products and lack the cytodifferentiation of mature islet cells. Pictet and Rutter, *supra*. Around day 16 in mouse gestation, the protodifferentiated pancreas begins a phase of rapid growth and differentiation characterized by cytodifferentiation of islet cells and a several hundred fold increase in islet specific gene expression. Histologically, islet formation (neogenesis) becomes apparent as proliferating islets bud from the pancreatic ducts (nesidioblastosis). Just before birth the rate of islet growth slows, and islet neogenesis and nesidioblastosis becomes much less apparent. Concomitant with this, the islets attain a fully differentiated state with maximal levels of insulin gene expression. Therefore, similar to many organs, the completion of cellular differentiation is associated with reduced regenerative potential; the differentiated adult pancreas does not have either the same regenerative potential or proliferative capacity as the developing pancreas.

Since differentiation of protodifferentiated precursors occurs during late fetal development of the pancreas, the factors regulating islet differentiation are likely to be expressed in the pancreas during this period. One of the genes expressed during islet

development encodes the gastrointestinal peptide, gastrin. Although gastrin acts in the adult as a gastric hormone regulating acid secretion, the major site of gastrin expression in the fetus is the pancreatic islets. Brand and Fuller, *J. Biol. Chem.*, 263:5341-5347 (1988). Expression of gastrin in the pancreatic islets is transient. It is confined to the period when protodifferentiated islet precursors form differentiated islets. Although the significance of pancreatic gastrin in islet development is unknown, some clinical observations suggest a role for gastrin in this islet development as follows. For example, hypergastrinemia caused by gastrin-expressing islet cell tumors and atrophic gastritis is associated with nesidioblastosis similar to that seen in differentiating fetal islets. Sacchi, *et al.*, *Virchows Archiv B*, 48:261-276 (1985); and Heitz *et al.*, *Diabetes*, 26:632-642 (1977). Further, an abnormal persistence of pancreatic gastrin has been documented in a case of infantile nesidioblastosis. Hollande, *et al.*, *Gastroenterology*, 71:251-262 (1976). However, in neither observation was a causal relationship established between the nesidioblastosis and gastrin stimulation.

It is therefore of interest to identify agents that stimulate islet cell regeneration which could be of value in the treatment of early IDDM and in the prevention of  $\beta$ -cell deficiency in NIDDM.

Citations of a reference herein shall not be construed as an admission that such reference is prior art to the present invention.

#### RELEVANT LITERATURE

Three growth factors are implicated in the development of the fetal pancreas, gastrin, transforming growth factor  $\alpha$  (TGF- $\alpha$ ) and epidermal growth factor (EGF) (Brand and Fuller, *J. Biol. Chem.* 263:5341-5347). Transgenic mice over expressing TGF- $\alpha$  or gastrin alone did not demonstrate active islet cell growth, however mice expressing both transgenes displayed significantly increased islet cell mass (Wang *et al.*, (1993) *J Clin Invest* 92:1349-1356). Bouwens and Pipeleers (1998) *Diabetologia* 41:629-633 report that there is a high proportion of budding  $\beta$ -cells in the normal adult human pancreas and 15% of all  $\beta$ -cells were found as single units. Single  $\beta$ -cell foci are not commonly seen in adult (unstimulated) rat pancreas; Wang *et al.* ((1995) *Diabetologia* 38:1405-1411) report a frequency of approximately 1% of total  $\beta$ -cell number.

Insulin independence in a Type 1 diabetic patient after encapsulated islet transplantation is described in Soon-Shiong *et al.* (1994) *Lancet* 343:950-51. Also see Sasaki

et al (1998 Jun 15) *Transplantation* 65(11):1510-1512; Zhou et al (1998 May) *Am J Physiol* 274(5 Pt 1):C1356-1362; Soon-Shiong et al (1990 Jun) *Postgrad Med* 87(8):133-134; Kendall et al (1996 Jun) *Diabetes Metab* 22(3):157-163; Sandler et al (1997 Jun) *Transplantation* 63(12):1712-1718; Suzuki et al (1998 Jan) *Cell Transplant* 7(1):47-52; 5 Soon-Shiong et al (1993 Jun) *Proc Natl Acad Sci USA* 90(12):5843-5847; Soon-Shiong et al (1992 Nov) *Transplantation* 54(5):769-774; Soon-Shiong et al (1992 Oct) *ASAIO J* 38(4):851-854; Benhamou et al (1998 Jun) *Diabetes Metab* 24(3):215-224; Christiansen et al (1994 Dec) *J Clin Endocrinol Metab* 79(6):1561-1569; Fraga et al (1998 Apr) *Transplantation* 65(8):1060-1066; Korsgren et al (1993) *Ups J Med Sci* 98(1):39-52; Newgard et al (1997 Jul) *Diabetologiz* 40 Suppl 2:S42-S47.

### SUMMARY OF THE INVENTION

The invention provides methods for treating diabetes mellitus in a patient in need thereof by administering a composition providing a gastrin/CCK receptor ligand, an EGF receptor ligand, or a combination of both in an amount sufficient to effect differentiation of the patient's pancreatic islet precursor cells to mature insulin-secreting cells. The composition can be administered systemically or expressed *in situ* by host cells containing a nucleic acid construct in an expression vector wherein the nucleic acid construct comprises a coding sequence for a gastrin CCK receptor ligand or a coding sequence for an EGF receptor ligand, together with transcriptional and translational regulatory regions functional in pancreatic islet precursor cells. Also provided are methods and compositions for treating diabetes in a patient in need thereof by implanting into a diabetic patient pancreatic islet cells that have been exposed in culture to a sufficient amount of a gastrin/CCK receptor ligand and an EGF receptor ligand to increase the number of pancreatic beta cells in the islets; 15 optionally the population of pancreatic beta cells can be grown in culture for a time sufficient to expand the population of  $\beta$ -cells prior to transplantation. The methods and compositions find use in treating patients with diabetes.

### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1A is an image that shows numerous insulin staining cells in the metaplastic ducts from the TGF- $\alpha$  transgenic pancreas upon immunoperoxidase staining. Figure 1B is an

image that shows that most ductular cells stained less intensely for insulin, while occasional ductular cells did stain with the same intensity of insulin staining as the adjacent islets.

Figure 2A schematically shows the structure of the chimeric insulin promoter-gastrin (INSGAS) transgene. Figure 2B illustrates that the radioimmunoassay of pancreatic extracts from INSGAS transgenic mice shows high levels of gastrin immunoreactivity that exceed the gastrin content in the gastric antrum expressed from the endogenous murine gene. The INSGAS transgenic mice had high expression of gastrin in the postnatal pancreas.

Figure 3A is an image of the pancreatic histology of an INSGAS/TGF- $\alpha$  mouse used in the study reported by Example 3. The INSGAS/TGF- $\alpha$  pancreas had some areas of increased ductular complexes and slightly increased interstitial cellularity. The field shown here had the most severely abnormal histology in the five animals used. Figure 3B is an image of the pancreatic histology of a control mouse from Example 3. Figure 3C is an image of the pancreatic histology of a TGF- $\alpha$  mouse from Example 3. This field of a TGF- $\alpha$  mouse pancreas from the study reported in Example 3 was typical and showed the interstitial cellularity and fibrosis combined with florid ductular metaplasia that has been described by Jhappan, *et al*, *supra*.

Figure 4A is a histogram graphically illustrating point-counting morphometric data which confirmed that at 17 weeks the pancreas of the INSGAS/TGF- $\alpha$  mice had lower duct mass than the pancreas of the TGF- $\alpha$  mice based on the study reported in Example 3. Figure 4B is a histogram which graphically illustrates point-counting morphometric data which show that co-expression of gastrin and TGF- $\alpha$  in the INSGAS/TGF- $\alpha$  pancreas significantly increased the islet mass compared to the islet mass of the corresponding non-transgenic control mice. Further, TGF- $\alpha$  expression alone does not increase islet mass. These data are based on the studies illustrated in Example 3.

Figure 5 shows the effects of TGF- $\alpha$  and gastrin on glucose tolerance in streptozotocin induced diabetic Wistar rats treated with PBS (solid black diamonds) or a combination of TGF- $\alpha$  and gastrin i.p. daily for 10 days (solid purple squares).

Figure 6 shows the effect of TGF- $\alpha$  and gastrin treatment on  $\beta$ -cell neogenesis in three groups of treated Zucker rats together with the corresponding PBS controls (n = 6 per group) as described in Example 7. The light blue bar represents lean TFG + gastrin, the magenta bar represents ob TGF + gastrin, the yellow bar represents the ob PBS control, the dark blue bar represents pre TFG + gastrin and the purple bar represents the lean PBS control.

TGF- $\alpha$  and gastrin significantly increased the relative proportion of single  $\beta$ -cell foci in all the groups studied as compared to PBS-treated control animals. Groups 4 and 5 are significantly different ( $p < 0.0015$ ) as are Groups 1 and 2 ( $p < 0.0041$ ).

Figure 7 shows the effect of TGF- $\alpha$  and gastrin treatment on  $\beta$ -cell neogenesis in lean and obese Zucker rats.  $\beta$ -cell neogenesis is quantified by differential counting of total  $\beta$ -cells and newly generated single  $\beta$ -cell foci and is expressed as a percentage of total  $\beta$ -cells counted. The percentage of single  $\beta$ -cell foci in lean Zucker rats treated with the growth factor combination was  $10.5 \pm 0.9$  compared to  $3.9 \pm 1.1$  ( $p = 0.004$ ) in the corresponding PBS control (Figs. 7A and 7B). In the obese Zucker rats, the percent single  $\beta$ -cell foci in the pretreatment group was  $8.7 \pm 1.3$  vs.  $4.2 \pm 1.1$  ( $p = 0.0015$ ) in the corresponding control group (Figs. 7C and 7D). Fig. 7E is a 400 x magnification of the ductal region of Fig. 7C (indicated by an arrow) and provides clear evidence of the budding of insulin-containing  $\beta$ -cells from the ductal epithelial cells characteristic of  $\beta$ -cell neogenesis.

#### **DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS**

The invention provides methods for treating diabetes mellitus in a patient in need thereof by administering a composition providing a gastrin/CCK receptor ligand such as gastrin, an EGF receptor ligand, such as TGF- $\alpha$ , or a combination of both in an amount sufficient to effect differentiation of pancreatic islet precursor cells to mature insulin-secreting cells. When the composition is administered systemically, generally it is provided by injection, preferably intravenously, in a physiologically acceptable carrier. When the composition is expressed *in situ*, pancreatic islet precursor cells are transformed either *in vivo* or *in vivo* with one or more nucleic acid expression constructs in an expression vector which provides for expression of the desired receptor ligand(s) in the pancreatic islet precursor cells. As an example, the expression construct includes a coding sequence for a CCK receptor ligand, such as preprogastrin peptide precursor coding sequence which, following expression, is processed to gastrin or a coding sequence for an EGF receptor ligand such as TGF- $\alpha$ , together with transcriptional and translational regulatory regions which provide for expression in the pancreatic islet precursor cells. The transcriptional regulatory region can be constitutive or induced, for example by increasing intracellular glucose concentrations, such as a transcriptional regulatory region from an insulin gene. Transformation is carried out using any suitable expression vector, for example, an adenoviral

expression vector. When the transformation is carried out *ex vivo*, the transformed cells are implanted in the diabetic patient, for example using a kidney capsule. Alternatively, pancreatic islet cells are treated *ex vivo* with a sufficient amount of a gastrin/CCK receptor ligand and/or an EGF receptor ligand to increase the number of pancreatic  $\beta$  cells in the islets prior to implantation into the diabetic patient. As required, the population of pancreatic  $\beta$  cells is expanded in culture prior to implantation by contacting them with the same receptor ligand(s).

The subject invention offers advantages over existing treatment regimens for diabetic patients. By providing a means to stimulate the adult pancreas to regenerate, not only is the need for traditional drug therapy (Type 2) or insulin therapy (Type 1 and Type 2) reduced or even eliminated, but the maintenance of normal blood glucose levels also may reduce some of the more debilitating complications of diabetes. Diabetic complications include those affecting the small blood vessels in the retina, kidney, and nerves, (microvascular complications), and those affecting the large blood vessels supplying the heart, brain, and lower limbs (macrovascular complications). Diabetic microvascular complications are the leading cause of new blindness in people 20-74 years old, and account for 35% of all new cases of end-stage renal disease. Over 60% of diabetics are affected by neuropathy. Diabetes accounts for 50% of all non-traumatic amputations in the USA, primarily as a result of diabetic macrovascular disease, and diabetics have a death rate from coronary artery disease that is 2.5 times that of non-diabetics. Hyperglycemia is believed to initiate and accelerate progression of diabetic microvascular complications. Use of the various current treatment regimens cannot adequately control hyperglycemia and therefore does not prevent or decrease progression of diabetic complications.

As used herein, the term "gastrin/CCK receptor ligand" encompasses compounds that stimulate the gastrin/CCK receptor. Examples of such gastrin/CCK receptor ligands include various forms of gastrin such as gastrin 34 (big gastrin), gastrin 17 (little gastrin), and gastrin 8 (mini gastrin); various forms of cholecystokinin such as CCK 58, CCK 33, CCK 22, CCK 12 and CCK 8; and other gastrin/CCK receptor ligands that either alone or in combination with EGF receptor ligands can induce differentiation of cells in mature pancreas to form insulin-secreting islet cells. Also contemplated are active analogs, fragments and other modifications of the above. Such ligands also include compounds that increase the secretion of endogenous gastrins, cholecystokinins or similarly active peptides from sites of tissue



storage. Examples of these are omeprazole which inhibits gastric acid secretion and soy bean trypsin inhibitor which increases CCK stimulation.

As used herein, the term "EGF receptor ligand" encompasses compounds that stimulate the EGF receptor such that when gastrin/CCK receptors in the same or adjacent tissues or in the same individual also are stimulated, neogenesis of insulin-producing pancreatic islet cells is induced. Examples of such EGF receptor ligands include EGF1-53, and fragments and active analogs thereof, including EGF1-48, EGF1-52, EGF1-49. See, for example, USPN 5,434,135. Other examples include TGF- $\alpha$  receptor ligands (1-50) and fragments and active analogs thereof, including 1-48, 1-47 and other EGF receptor ligands such as amphiregulin and pox virus growth factor as well as other EGF receptor ligands that demonstrate the same synergistic activity with gastrin/CCK receptor ligands. These include active analogs, fragments and modifications of the above. For further background, see Carpenter and Wahl, Chapter 4 in *Peptide Growth Factors* (Eds. Sporn and Roberts), Springer Verlag, (1990).

A principal aspect of the invention is a method for treating diabetes mellitus in an individual in need thereof by administering to the individual a composition including a gastrin/CCK receptor ligand and/or an EGF receptor ligand in an amount sufficient to effect differentiation of pancreatic islet precursor cells to mature insulin-secreting cells. The cells so differentiated are residual latent islet precursor cells in the pancreatic duct. One embodiment comprises administering, preferably systemically, a differentiation regenerative amount of a gastrin/CCK receptor ligand and an EGF receptor ligand, preferably TGF- $\alpha$ , either alone or in combination to the individual.

Another embodiment comprises providing a gastrin/CCK receptor ligand and/or an EGF receptor ligand to pancreatic islet precursor cells of explanted pancreatic tissue of a mammal and reintroducing the pancreatic tissue so stimulated to the mammal.

In another, the invention comprises providing a gastrin/CCK receptor ligand and/or an EGF receptor ligand to pancreatic islet precursor cells of explanted pancreatic tissue from a mammal to expand the population of  $\beta$  cells.

In another embodiment gastrin/CCK receptor ligand stimulation is effected by expression of a chimeric insulin promoter-gastrin fusion gene construct transgenically introduced into such precursor cells. In another embodiment EGF receptor ligand stimulation

is effected by expression of an EGF receptor ligand gene transgenically introduced into the mammal. The sequence of the EGF gene is provided in USPN 5,434,135.

In another embodiment stimulation by a gastrin/CCK receptor ligand and an EGF receptor ligand is effected by coexpression of (i) a preprogastrin peptide precursor gene and (ii) an EGF receptor ligand gene that have been stably introduced into the mammal.

In another aspect the invention relates to a method for effecting the differentiation of pancreatic islet precursor cells of a mammal by stimulating such cells with a combination of a gastrin/CCK receptor ligand and an EGF receptor ligand. In a preferred embodiment of this aspect, gastrin stimulation is effected by expression of a preprogastrin peptide precursor gene stably introduced into the mammal. The expression is under the control of the insulin promoter. EGF receptor ligand, e.g., TGF- $\alpha$ , stimulation is effected by expression of an EGF receptor ligand gene transgenically introduced into the mammal. In furtherance of the above, stimulation by a gastrin and a TGF- $\alpha$  is preferably effected by co-expression of (i) a preprogastrin peptide precursor gene and (ii) an EGF receptor ligand introduced into the mammal. Appropriate promoters capable of directing transcription of the genes include both viral promoters and cellular promoters. Viral promoters include the immediate early cytomegalovirus (CMV) promoter (Boshart *et al* (1985) *Cell* 41:521-530), the SV40 promoter (Subramani *et al* (1981) *Mol. Cell. Biol.* 1:854-864) and the major late promoter from Adenovirus 2 (Kaufman and Sharp (1982) *Mol. Cell. Biol.* 2:1304-1319). Preferably, expression of one or both of the gastrin/CCK receptor ligand gene and the EGF receptor ligand gene is under the control of an insulin promoter.

Another aspect of the invention is a nucleic acid construct. This construct includes a nucleic acid sequence coding for a preprogastrin peptide precursor and an insulin transcriptional regulatory sequence, which is 5' to and effective to support transcription of a sequence encoding the preprogastrin peptide precursor. Preferably, the insulin transcriptional regulatory sequence includes at least an insulin promoter. In a preferred embodiment the nucleic acid sequence coding for the preprogastrin peptide precursor comprises a polynucleotide sequence containing exons 2 and 3 of a human gastrin gene and optionally also including introns 1 and 2.

Another embodiment of the invention is an expression cassette comprising (i) a nucleic acid sequence coding for a mammalian EGF receptor ligand, e.g., TGF- $\alpha$  and a transcriptional regulatory sequence thereof; and (ii) a nucleic acid sequence coding for the

preprogastrin peptide precursor and a transcriptional regulatory sequence thereof. Preferably, the transcriptional regulatory sequence for the EGF receptor ligand is a strong non-tissue specific promoter, such as a metallothionein promoter. Preferably, the transcriptional regulatory sequence for the preprogastrin peptide precursor is an insulin promoter. A preferred form of this embodiment is one wherein the nucleic acid sequence coding for the preprogastrin peptide precursor comprises a polynucleotide sequence containing introns 1 and 2 and exons 2 and 3 of the human gastrin gene.

Another aspect of the invention relates to a vector including the expression cassette comprising the preprogastrin peptide precursor coding sequence. This vector can be a plasmid such as pGem1 or can be a phage which has a transcriptional regulatory sequence including an insulin promoter.

Another aspect of this invention relates to a composition of vectors including (1) having the nucleic acid sequence coding for a mammalian EGF receptor ligand, e.g., TGF- $\alpha$ , under control of a strong non-tissue specific promoter, e.g., a metallothionein promoter; and a preprogastrin peptide precursor coding sequence under control of an insulin promoter. Each vector can be a plasmid, such as plasmid pGem1 or a phage in this aspect. Alternatively, the expression cassette or vector also can be inserted into a viral vector with the appropriate tissue tropism. Examples of viral vectors include adenovirus, *Herpes simplex* virus, adeno-associated virus, retrovirus, lentivirus, and the like. See Blomer *et al* (1996) *Human Molecular Genetics* 5 Spec. No:1397-404; and Robbins *et al* (1998) *Trends in Biotechnology* 16:35-40. Adenovirus-mediated gene therapy has been used successfully to transiently correct the chloride transport defect in nasal epithelia of patients with cystic fibrosis. See Zabner *et al.* (1993) *Cell* 75:207-216.

Another aspect of the invention is a non-human mammal or mammalian tissue, including cells, thereof capable of expressing a stably integrated gene which encodes preprogastrin. Another embodiment of this aspect is a non-human mammal capable of coexpressing (i) a preprogastrin peptide precursor gene; and/or (ii) an EGF receptor ligand, e.g., a TGF- $\alpha$  gene that has been stably integrated into the non-human mammal, mammalian tissue or cells. The mammalian tissue or cells can be human tissue or cells.

### **Therapeutic Administration and Compositions**

Modes of administration include but are not limited to transdermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, and oral routes. The compounds can be administered by any convenient route, for example by infusion or bolus injection by absorption through epithelial or mucocutaneous linings (e.g., oral mucosa, rectal and intestinal mucosa, etc.) and can be administered together with other biologically active agents. Administration is preferably systemic.

The present invention also provides pharmaceutical compositions. Such compositions comprise a therapeutically effective amount of a therapeutic, and a pharmaceutically acceptable carrier or excipient. Such a carrier includes but is not limited to saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. The formulation should suit the mode of administration. Pharmaceutically acceptable carriers and formulations for use in the present invention are found in Remington's *Pharmaceutical Sciences*, Mack Publishing Company, Philadelphia, PA, 17<sup>th</sup> ed. (1985), which is incorporated herein by reference. For a brief review of methods for drug delivery, see Langer (1990) *Science* 249:1527-1533, which is incorporated herein by reference.

In preparing pharmaceutical compositions of the present invention, it may be desirable to modify the compositions of the present invention to alter their pharmacokinetics and biodistribution. For a general discussion of pharmacokinetics, see Remington's *Pharmaceutical Sciences*, *supra*, Chapters 37-39. A number of methods for altering pharmacokinetics and biodistribution are known to one of ordinary skill in the art (See, e.g., Langer, *supra*). Examples of such methods include protection of the agents in vesicles composed of substances such as proteins, lipids (for example, liposomes), carbohydrates, or synthetic polymers. For example, the agents of the present invention can be incorporated into liposomes in order to enhance their pharmacokinetics and biodistribution characteristics. A variety of methods are available for preparing liposomes, as described in, e.g., Szoka *et al* (1980) *Ann. Rev. Biophys. Bioeng.* 9:467, U.S. Pat. Nos. 4,235,871, 4,501,728 and 4,837,028, all of which are incorporated herein by reference. Various other delivery systems are known and can be used to administer a therapeutic of the invention, e.g., microparticles, microcapsules and the like.

The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. The composition can be a liquid solution,

suspension, emulsion, tablet, pill, capsule, sustained release formulation, or powder. The composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. Oral formulations can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, etc.

In a preferred embodiment, the composition is formulated in accordance with routine procedures such as a pharmaceutical composition adapted for intravenous administration to human beings. Typically, compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. Where necessary, the composition also can include a solubilizing agent and a local anesthetic to ameliorate any pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

The therapeutics of the invention can be formulated as neutral or salt forms. Pharmaceutically acceptable salts include those formed with free amino groups such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., and those formed with free carboxyl groups such as those derived from sodium, potassium, ammonium, calcium and other divalent cations, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc.

The amount of the therapeutic of the invention which is effective in the treatment of a particular disorder or condition will depend on the nature of the disorder or condition, and can be determined by standard clinical techniques. The precise dose to be employed in the formulation also will depend on the route of administration, and the seriousness of the disease or disorder, and should be decided according to the judgment of the practitioner and each patient's circumstances. However, suitable dosage ranges for intravenous administration are generally about 20-500 micrograms of active compound per kilogram body weight. Suitable dosage ranges for intranasal administration are generally about 0.01 pg/kg body weight to 1 mg/kg body weight. Effective dosages can be extrapolated from dose-response curves

derived from *in vitro* or animal model test systems. Suppositories generally contain active ingredient in the range of 0.5% to 10% weight; oral formulations preferably contain 10% to 95% active ingredient.

In the gene therapy methods of the invention, transfection *in vivo* is obtained by introducing a therapeutic transcription or expression vector into the mammalian host, either as naked DNA, complexed to lipid carriers, particularly cationic lipid carriers, or inserted into a viral vector, for example a recombinant adenovirus. The introduction into the mammalian host can be by any of several routes, including intravenous or intraperitoneal injection, intratracheally, intrathecally, parenterally, intraarticularly, intranasally, intramuscularly, topically, transdermally, application to any mucous membrane surface, corneal installation, etc. Of particular interest is the introduction of the therapeutic expression vector into a circulating bodily fluid or into a body orifice or cavity. Thus, intravenous administration and intrathecal administration are of particular interest since the vector may be widely disseminated following such routes of administration, and aerosol administration finds use with introduction into a body orifice or cavity. Particular cells and tissues can be targeted, depending upon the route of administration and the site of administration. For example, a tissue which is closest to the site of injection in the direction of blood flow can be transfected in the absence of any specific targeting. If lipid carriers are used, they can be modified to direct the complexes to particular types of cells using site-directing molecules. Thus, antibodies or ligands for particular receptors or other cell surface proteins may be employed, with a target cell associated with a particular surface protein.

Any physiologically acceptable medium may be employed for administering the DNA, recombinant viral vectors or lipid carriers, such as deionized water, saline, phosphate-buffered saline, 5% dextrose in water, and the like as described above for the pharmaceutical composition, depending upon the route of administration. Other components can be included in the formulation such as buffers, stabilizers, biocides, etc. These components have found extensive exemplification in the literature and need not be described in particular here. Any diluent or components of diluents that would cause aggregation of the complexes should be avoided, including high salt, chelating agents, and the like.

The amount of therapeutic vector used will be an amount sufficient to provide for a therapeutic level of expression in a target tissue. A therapeutic level of expression is a sufficient amount of expression to decrease blood glucose towards normal levels. In addition,

the dose of the nucleic acid vector used must be sufficient to produce a desired level of transgene expression in the affected tissues *in vivo*. Other DNA sequences, such as adenovirus VA genes can be included in the administration medium and be co-transfected with the gene of interest. The presence of genes coding for the adenovirus VA gene product may significantly enhance the translation of mRNA transcribed from the expression cassette if this is desired.

A number of factors can affect the amount of expression in transfected tissue and thus can be used to modify the level of expression to fit a particular purpose. Where a high level of expression is desired, all factors can be optimized, where less expression is desired, one or more parameters can be altered so that the desired level of expression is attained. For example, if high expression would exceed the therapeutic window, then less than optimum conditions can be used.

The level and tissues of expression of the recombinant gene may be determined at the mRNA level as described above, and/or at the level of polypeptide or protein. Gene product may be quantitated by measuring its biological activity in tissues. For example, protein activity can be measured by immunoassay as described above, by biological assay such as blood glucose, or by identifying the gene product in transfected cells by immunostaining techniques such as probing with an antibody which specifically recognizes the gene product or a reporter gene product present in the expression cassette.

Typically, the therapeutic cassette is not integrated into the patient's genome. If necessary, the treatment can be repeated on an *ad hoc* basis depending upon the results achieved. If the treatment is repeated, the patient can be monitored to ensure that there is no adverse immune or other response to the treatment.

The invention also provides for methods for expanding a population of pancreatic  $\beta$ -cells *in vitro*. Upon isolation of the pancreas from a suitable donor, cells are isolated and grown *in vitro*. The cells which are employed are obtained from tissue samples from mammalian donors including human cadavers, porcine fetuses or another suitable source of pancreatic cells. If human cells are used, when possible the cells are major histocompatibility matched with the recipient. Purification of the cells can be accomplished by gradient separation after enzymatic (e.g., collagenase) digestion of the isolated pancreas. The purified cells are grown in media containing sufficient nutrients to allow for survival of the cells as well as a sufficient amount of a  $\beta$ -cell proliferation inducing composition

containing a gastrin/CCK receptor ligand and EGF receptor ligand, to allow for formation of insulin secreting pancreatic  $\beta$  cells. According to the invention, following stimulation the insulin secreting pancreatic  $\beta$  cells can be directly expanded in culture prior to being transplanted into a patient in need thereof, or can be transplanted directly following treatment with  $\beta$ -cell proliferation inducing composition.

Methods of transplantation include transplanting insulin secreting pancreatic  $\beta$ -cells obtained into a patient in need thereof in combination with immunosuppressive agents, such as cyclosporin. The insulin producing cells also can be encapsulated in a semi-permeable membrane prior to transplantation. Such membranes permit insulin secretion from the encapsulated cells while protecting the cells from immune attack. The number of cells to be transplanted is estimated to be between 10,000 and 20,000 insulin producing  $\beta$  cells per kg of the patient. Repeated transplants may be required as necessary to maintain an effective therapeutic number of insulin secreting cells. The transplant recipient can also, according to the invention, be provided with a sufficient amount of a gastrin/CCK receptor ligand and an EGF receptor ligand to induce proliferation of the transplanted insulin secreting  $\beta$  cells.

The effect of treatment of diabetes can be evaluated as follows. Both the biological efficacy of the treatment modality as well as the clinical efficacy are evaluated, if possible. For example, disease manifests itself by increased blood sugar, the biological efficacy of the treatment therefore can be evaluated, for example, by observation of return of the evaluated blood glucose towards normal. The clinical efficacy, i.e. whether treatment of the underlying effect is effective in changing the course of disease, can be more difficult to measure. While the evaluation of the biological efficacy goes a long way as a surrogate endpoint for the clinical efficacy, it is not definitive. Thus, measuring a clinical endpoint which can give an indication of  $\beta$ -cell regeneration after, for example, a six-month period of time, can give an indication of the clinical efficacy of the treatment regimen.

The subject compositions can be provided as kits for use in one or more procedures. Kits for genetic therapy usually will include the therapeutic DNA construct either as naked DNA with or without mitochondrial targeting sequence peptides, as a recombinant viral vector or complexed to lipid carriers. Additionally, lipid carriers can be provided in separate containers for complexing with the provided DNA. The kits include a composition comprising an effective agent either as concentrates (including lyophilized compositions), which can be diluted further prior to use or they can be provided at the concentration of use,



where the vials may include one or more dosages. Conveniently, in the kits single dosages can be provided in sterile vials so that the physician can employ the vials directly, where the vials will have the desired amount and concentration of agents. When the vials contain the formulation for direct use, usually there will be no need for other reagents for use with the method. Associated with such kits can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

The following examples are offered by way of illustration and not by way of limitation.

### **EXAMPLES**

#### **Materials and Methods**

The following materials and methods were used in the studies reported by the working examples set forth below except as otherwise noted.

**Animals.** Mice, FVB and CD strain, were obtained from Taconic Farms, Inc., Germantown, NY. The TGF- $\alpha$  transgenic line MT-42 used, which expresses high levels of TGF- $\alpha$  from a metallothionein promoter, is described in Jhappan *et al*, *Cell*, 61:1137-1146 (1990). Normal Wistar and Zucker rats were allowed normal chow *ad libidum* with free access to water and were acclimatized for one week prior to initiation of each study. Freshly prepared streptozotocin at a dose of 80 mg/kg body weight was administered by I.V. five to seven days after induction of diabetes, the rats were randomly allocated into groups for subsequent treatment. Hormones, TGF- $\alpha$  and rat gastrin were reconstituted in sterile normal saline containing 0.1% BSA. According to the predetermined treatment schedule for different studies, each animal received a single, daily i.p. injection of either TGF- $\alpha$  or gastrin alone (4.0  $\mu$ g/kg body weight) or as a 1:1 (w/w) combination (total 8.0  $\mu$ g/kg) or PBS for a period of 10 days.

**INS GAS Transgene Construct.** A *PvuII-RsaI* fragment encompassing nucleotides -370 to +38 of the rat insulin I gene (Cordell, B.G. *et al*, *Cell*, 18:533-543 (1979)) was ligated into pGem1 (Promega Corp., Madison, WI). A 4.4 kb *BamHI-EcoRI* fragment containing 1.5 kb introns

1 and 2 and exons 2 and 3 of the human gastrin gene which encodes the preprogastrin peptide precursor was isolated and subcloned downstream of the rat insulin I fragment in pGem1 (Promega). The fragment is described in Wiborg, O., *Proc. Natl. Acad. Sci. USA*, 81:1067-1069 (1984) and Ito, R., *et al Proc. Natl. Acad. Sci. (USA)*, 81:4662-4666 (1984). The insulin promoter-preprogastrin INSGAS transgene construct was excised as a 4.8 kb *Xba*I-*Eco*R1 fragment.

Generation and Characterization of Transgenic Mice. The fragment, made as described above was prepared for microinjection as follows. It was isolated by agarose gel electrophoresis, purified by CsCl gradient purification, and dialyzed extensively against injection buffer (5mM NaCl; 0.1 mM EDTA; 5mM Tris-HCl pH 7.4). Fertilized oocytes from FVB inbred mice (Taconic Farms, Inc., *supra*) at the single-cell stage were microinjected using standard techniques. See Hogan, B., *et al, Manipulating the mouse embryo: A laboratory manual*, Cold Spring Harbor, NY (1986). Surviving embryos were then implanted into the oviducts of CD1 (Charles River Laboratories, Inc., Wilmington, MA) foster mothers according to procedures in Hogan *et al*. Transgenic founder mice were identified by DNA blot techniques using DNA isolated from individual mouse tails, and a human gastrin exon 2 probe labelled with <sup>32</sup>P by random priming. F1 mice and their siblings were similarly identified.

Homozygous MT-42 mice containing the MT-TGF- $\alpha$  transgene derived from a CD-1 mouse strain (Jhappan, *supra*) were crossed with heterozygotic INSGAS mice. After weaning, the offspring were placed on acidified 50mM ZnCl<sub>2</sub> as previously described in order to induce the metallothionein promoter (Jhappan, *supra*).

Northern Blot Hybridization Assay. For Northern analysis, total RNA was extracted from tissues by the method of Cathala *et al*, *DNA* 2:329-335 (1983). Samples of 20 $\mu$ g of total RNA were resolved on a 1% agarose denaturing gel and transferred to nitrocellulose. RNA blots were hybridized with <sup>32</sup>P labelled TGF- $\alpha$  riboprobes or exon 2 of human gastrin that did not cross-hybridize with endogenous mouse gastrin mRNA.

Peptide radioimmunoassay of Gastrin. Tissues were extracted and assayed for gastrin immunoreactivity by radioimmunoassay as described previously using antibody 2604 which

is specific for biologically active C terminally amidated gastrin in a gastrin radioimmunoassay as described in Rehfeld, J.F., *Scand. J. Clin. Lab. Invest.* 30:361-368 (1972). Tyrosine moniodinated human gastrin 17 tracer was used in all assays and synthetic human gastrin 17 was used as a standard.

5

Peptide Radioimmunoassay of TGF- $\alpha$ : Tissues were frozen in liquid nitrogen, ground to a powder with mortar and pestle, and subjected to acid-ethanol extraction as described in Todaro, G.J. *et al*, *Proc. Natl. Acad. Sci. (USA)*, 77:5258-5262 (1980). Extracts were reconstituted with water, and protein concentrations determined with a Coomassie blue dye binding assay (Bio-Rad Laboratories, Hercules, CA). Aliquots from the pancreata were tested in duplicate in a TGF- $\alpha$  radioimmunoassay, which measured competition with  $^{125}\text{I}$  TGF- $\alpha$  for binding to a solid-phase rabbit antibody raised against the C-terminus of rat TGF- $\alpha$  (kit from BioTope, Seattle, WA).

10

Blood Glucose. Blood glucose was determined either after overnight fasting or after IPGTT by glucose oxidase method.

15

Tissue Insulin Analysis. At the end of each study, the animals were sacrificed and pancreas removed. Small biopsies were taken from separate representative sites throughout the pancreas and immediately snap-frozen in liquid nitrogen for immunohistochemistry, protein, and insulin determinations. Snap-frozen pancreatic samples ( $n = 5$ ) were rapidly thawed, disrupted ultrasonically in deionized water and aliquots taken for protein determination and the homogenate subjected to acid/ethanol extraction prior to insulin determination by RIA.

20

Histological Analysis. The pancreata were removed, weighed, similarly oriented in cassettes, fixed in Bouin's solution and embedded in paraffin by conventional procedures.

25

Tissue Preparation and Immunohistochemistry. Freshly excised pancreata were dissected, cleared of fat and lymph nodes, fixed in Bouin's fixative, and then embedded in paraffin for sectioning. Routine sections were stained with hematoxylin and eosin according to standard methods. Pancreatic tissue from adult 17 week old MT-TGF- $\alpha$  (MT-42) transgenic mice were immunostained for insulin to examine the effect of TGF- $\alpha$  over-expression on islet

30

development. Insulin positive cells in TGF- $\alpha$ -induced metaplastic ductules were identified using immunoperoxidase staining guinea pig anti-human insulin sera (Linco, Eureka, MO); a pre-immune guinea pig serum was used as a control. Immunohistochemistry was performed on 5 $\mu$  paraffin sections by the peroxidase/antiperoxidase method of Sternberger using a monoclonal rabbit antigastrin antibody. See, Sternberger, L.A., *Immunocytochemistry*, 2<sup>nd</sup> Ed. (1979) NY: Wiley. 104-170.

Point-Counting Morphometrics. The relative volume of islets, ducts, or interstitial cells was quantitated using the point-counting method described in Weibel, E.R., *Lab Invest.*, 12:131-155 (1963). At a magnification of 400x, starting at a random point at one corner of the section, every other field was scored using a 25 point ocular grid. An unbiased but systematic selection of fields was accomplished using the markings of the stage micrometer. Intercepts over blood vessels, fat, ducts, lymph nodes, or interlobular space were subtracted to give the total pancreatic area. A minimum of 5000 points in 108 fields (systematically chosen using the stage micrometer) were counted in each block, with the relative islet volume being the number of intercepts over islet tissue divided by the number over pancreatic tissue. The absolute islet mass or islets was calculated as the relative islet volume times pancreatic weight. See, Lee, H.C., *et al*, *Endocrinology*, 124:1571-1575 (1989).

Statistical Analysis. Differences between means were compared for significant differences using the Student's *t* test for unpaired data.

### EXAMPLE 1

#### Assay For Insulin Production in TGF Transgenic Pancreas

Immunoperoxidase staining showed numerous insulin staining cells in the metaplastic ducts from the TGF- $\alpha$  transgenic pancreas (Figure 1A), whereas insulin staining cells were virtually absent from the non-transgenic ducts (less than 6.1%). When at least 600 ductular cells/animal were scored at a final magnification of 400x, insulin positive cells were seen at a frequency of 6.0 +/- 0.9% (n = 5) in the metaplastic ductules of TGF- $\alpha$  transgenic mice. Occasional ductular cells stained with the same intensity of insulin staining as the adjacent islets, but most had less intense staining (Figure 1B). The low level of insulin staining of the ductular cells resembles that of protodifferentiated cells reported in the ducts of the

developing pancreas. Pictet, R. and W.J. Rutter, *Development of the embryonic endocrine pancreas, in Endocrinology, Handbook of Physiology*, ed. R.O. Greep and E.B. Astwood (1972) American Physiological Society: Washington, D.C. 25-66; and Alpert, S. *et al Cell*, 53:295-308 (1988).

5           However, despite the increased number of insulin positive cells in the metaplastic ducts, the islet mass of the TGF- $\alpha$  transgenic mice was not increased. The islet mass as quantitated by point counting morphometrics was 2.14 mg  $\pm$  0.84 (mean  $\pm$  se, n = 5) in the TGF- $\alpha$  transgenic pancreas compared to 1.93 mg  $\pm$  0.46 (n = 6) non transgenic litter mates.

10           Thus, TGF- $\alpha$  over-expression alone did not effect transition of these protodifferentiated duct cells into fully differentiated islets. This implies that islet differentiation requires other factors absent from the adult pancreas of TGF- $\alpha$  transgenic mice. Since differentiation of protodifferentiated islet precursors occurs during late fetal development, factors regulating this transition would likely be expressed in islets during this period. Among the factors expressed in the developing islets are the gastrointestinal peptides, the gastrins.

## EXAMPLE 2

### Pancreatic Gastrin Expression from the INSGAS Transgene

15           To examine the possible role of gastrin in regulating islet differentiation, transgenic mice were created that express a chimeric insulin promoter-gastrin (INSGAS) transgene in which the insulin promoter directs pancreas specific expression of the gastrin transgene (Figure 2A). Unlike the gastrin gene, insulin gene expression is not switched off after birth. Thus, the INSGAS transgene results in a persistence of gastrin expression in the adult pancreas.

20           The INSGAS transgene comprised 370 bp of 5' flanking DNA and the first non-coding exon of the rat insulin I gene. Cordell, B., *et al, Cell* 18:533-543 (1979). It was ligated to a *Bam*H1-*Eco*R1 fragment containing 1.5 kb intron 1 and exons 2 and 3 of the human gastrin gene which encodes the preprogastrin peptide precursor. Wiborg, O., *et al, Proc. Natl. Acad. Sci. USA*, 81:1067-1069 (1984); and Ito *et al Proc. Natl. Acad. Sci. USA*, 81:4662-4666 (1984). A 4.8 kb INSGAS fragment was isolated and microinjected into inbred FVB, one cell mouse embryos. Hogan, B. *et al, Manipulating the mouse embryo: A laboratory manual*, (1986) NY: Cold Spring Harbor.

Gastrin immunoreactivity in pancreatic and stomach extracts from transgenic and non-transgenic mice was assayed by radioimmunoassay using antisera 2604 (Rehfeld, J., *et al*, *Scand. J. Clin. Lab. Invest.*, 30:361-368 (1972)) specific for the bioactive amidated C-terminus of gastrin.

5       Beta cell specific gastrin expression from the INSGAS transgene was observed based on immunostaining of pancreatic tissues with a gastrin monoclonal antibody.

Northern blots of RNA isolated from different tissues of 8 week old INSGAS transgenic mice were hybridized with a human gastrin exon 2 probe. High levels of gastrin transgene mRNA were seen in the pancreas but not in any other tissues. This probe is specific for the human gastrin gene; no hybridization is seen in antral RNA of INSGAS and non-transgenic FVB mice express high levels of murine gastrin mRNA. Radioimmunoassay of pancreatic extracts from INSGAS transgenic mice showed high levels of gastrin immunoreactivity that exceed the gastrin content in the gastric atrium expressed from the endogenous murine gene (Figure 2B). No gastrin immunoreactivity was detected in pancreatic extracts of non-transgenic control mice (Figure 2B). The gastrin radioimmunoassay is specific for carboxy amidated precursors, indicating that the gastrin peptide precursor is efficiently processed post-translationally to the bioactive peptide. Immunohistochemistry with a gastrin monoclonal antibody shows pancreatic beta islet cell specific expression of gastrin.

10  
15  
20       Although the INSGAS transgenic mice had high expression of gastrin in the postnatal pancreas (Figure 2B), the INSGAS transgenic mice had pancreatic histology identical to controls. Islet mass as quantitated by point-counting morphometrics (Weibel, E.R., *Lab Invest.* 12:131-155 (1963)) was identical in 5-6 week old INSGAS mice (1.78 +/- 0.21 mg, n = 11) and age matched non-transgenic controls (1.74 +/- 0.18mg, n = 11). Thus, sustained  
25       expression of gastrin in the postnatal pancreas alone does not stimulate islet cell growth.

### EXAMPLE 3

#### Histological Examination of TGF- $\alpha$ and TGF- $\alpha$ /INSGAS Pancreas

30       Stimulation of islet growth by gastrin may require stimulation by other growth factors to create a responsive population of cells. Therefore, effects of gastrin stimulation were studied in TGF- $\alpha$  transgenic mice which have metaplastic ducts that contain insulin expressing cells resembling protodifferentiated islet-precursors. To assess the interaction

between gastrin and TGF- $\alpha$ , three groups of mice were bred with equivalent FVB/CD1 strain genetic backgrounds: non-transgenic control, TGF- $\alpha$  single transgenic and INSGAS/TGF- $\alpha$  double transgenics. All three groups of mice were placed on 50mM ZnCl<sub>2</sub> at 3 weeks of age.

At 17 weeks of age, the animals were sacrificed and the pancreas removed for histological evaluation. The pancreas from TGF- $\alpha$  and INSGAS/TGF- $\alpha$  mice had similar gross morphological appearances: resilient, firm and compact in contrast to the soft diffuse control pancreas. TGF- $\alpha$  expression was equivalent in TGF- $\alpha$  and INSGAS/TGF- $\alpha$  groups when measured by Northern blot analysis (data not shown) and by radioimmunoassay. The pancreatic TGF- $\alpha$  immunoreactive peptide levels were 12.2 +/- 1 and 18.9 +/- 8ng/mg protein (Mean +/- SD) in the TGF- $\alpha$  and INSGAS/TGF- $\alpha$  mice, respectively.

Light micrographs of hematoxylin stained paraffin sections of pancreas from the three groups of mice studied (A: INSGAS/TGF- $\alpha$ ; B: FVB/CD1 controls; and C: TGF- $\alpha$ ) were made. The INSGAS/TGF- $\alpha$  pancreas had some areas of increased ductular complexes and slightly increased interstitial cellularity; the field shown (Figure 3A) had the most severely abnormal morphology seen in the five animals; most of the pancreas was indistinguishable from controls (Figure 3B). In contrast, the field of TGF- $\alpha$  pancreas (Figure 3C) was typical and showed the interstitial cellularity and fibrosis combined with florid ductular metaplasia described by Jhappan *et al*, *supra*.

Pancreatic gastrin synergistically interacts with TGF- $\alpha$  to increase islet mass and inhibit the ductular metaplasia induced by TGF- $\alpha$  over-expression. Mating the homozygous MT-TGF- $\alpha$  (MT-42) mice (TGF- $\alpha$ ) with heterozygotic INSGAS mice gave offspring that were either heterozygotic TGF- $\alpha$  single transgenic or double transgenic containing both INSGAS and TGF- $\alpha$  transgenes (INSGAS/TGF- $\alpha$ ). Since INSGAS were FVB strain and TGF- $\alpha$  were CD1 strain, TGF- $\alpha$  homozygotes and CD1 controls (CON) were both mated with FVB to produce FVB/CD1 strain background for all three groups of mice. Mice were treated with 50mM ZnCl<sub>2</sub> from 3 weeks until sacrifice at age 17 weeks. The pancreas was removed, weighed, similarly oriented in cassettes, fixed in Bouin's solution and embedded in paraffin. One random section from each animal was used to quantitate the relative volumes of ductules and islets by point-counting morphometrics (Weibel, E.R., *Lab Invest.*, 12:131-155 (1963)). At least 2000 points over tissue were counted as intercepts of a 50 point grid at 170x magnification; the entire section was covered without overlap. The mass of ductules or islets was calculated by multiplying the relative volume and the animal's pancreatic weight.

To normalize different mean body weights, the mass was expressed as  $\mu\text{g/g}$  body weight. Results are mean and standard errors for 5-6 animals in each group as determined by Student's *t* test ( $p < 0.05$ ).

Expression of gastrin from the INSGAS transgene reduced the ductular metaplasia caused by TGF- $\alpha$  over-expression. At 17 weeks, the pancreatic histology of the INSGAS/TGF- $\alpha$  mice (Figure 3A) resembled that of the control pancreas (Figure 3B) more than that of the TGF- $\alpha$  mice (Figure 3C).

This was confirmed by quantitating pancreatic ductular mass in the TGF- $\alpha$  and INSGAS/TGF- $\alpha$  transgenic mice and the FVB/CD1 controls by point-counting morphometrics (Figure 4A). Co-expression of gastrin and TGF- $\alpha$  in the INSGAS/TGF- $\alpha$  pancreas also significantly increased the islet mass compared to controls (Figure 4B), whereas islet mass was not increased by expression of the TGF- $\alpha$  or gastrin transgenes alone. The blood glucose concentration was not significantly different among the three groups of mice.

#### EXAMPLE 4

##### Effects of TGF- $\alpha$ and Gastrin on Pancreatic Insulin Content in Normal Rats

This experiment was designed to study the effects on pancreatic insulin content in non-diabetic animals treated with TGF- $\alpha$ , a gastrin, or a combination of TGF- $\alpha$  and a gastrin as compared to control animals (untreated). Groups ( $n = 5$ ) of normal Wistar rats were assigned to one of the following four treatment groups.

Group I: TGF- $\alpha$ : recombinant Human TGF- $\alpha$  was reconstituted in sterile saline containing 0.1% BSA and was administered i.p. at a dose of 0.8  $\mu\text{g/day}$  for 10 days.

Group II: Gastrin: synthetic Rat Gastrin I was dissolved in very dilute ammonium hydroxide and reconstituted in sterile saline containing 0.1% BSA. It was administered i.p. at a dose of 0.8  $\mu\text{g/day}$  for 10 days.

Group III: TGF- $\alpha$  + Gastrin: a combination of the above preparations was administered i.p. at the dose levels given above for 10 days.



Group IV: Control animals received an i.p. injection of vehicle alone for 10 days.

At the end of the study period (10 days), all animals were sacrificed and samples of pancreas taken as follows: five biopsy specimens (1-2 mg) of pancreatic tissue were taken from separate representative sites in each rat pancreas and immediately snap frozen in liquid nitrogen for analysis of insulin content. For analysis of pancreatic insulin content, the snap frozen pancreatic samples were rapidly thawed, disrupted ultrasonically in distilled water and aliquots taken for protein determination and acid/ethanol extraction prior to insulin radioimmunoassay (Green *et al*, (1983) *Diabetes* 32:685-690). Pancreatic insulin content values were corrected according to protein content and finally expressed as  $\mu\text{g}$  insulin/mg pancreatic protein. All values calculated as mean  $\pm$  SEM and statistical significance evaluated using Student's 2-sample *t*-test.

**Table 1**

**Treatment of Normal Rats with TGF- $\alpha$  and Gastrin**

<b>Treatment</b>	<b><u>Pancreatic Insulin Content</u> (<math>\mu\text{g}</math> insulin/mg protein)</b>
Control	20.6 $\pm$ 6.0
TGF- $\alpha$	30.4 $\pm$ 7.4*
Gastrin	51.4 $\pm$ 14.0**
TGF- $\alpha$ + Gastrin	60.6 $\pm$ 8.7***

\* TGF- $\alpha$  vs. control,  $p = 0.34$ ;

\*\* gastrin vs. control,  $p = 0.11$ ;

\*\*\* combination of TGF- $\alpha$  and gastrin,  $p = 0.007$ .

As shown in Table 1, above, pancreatic insulin content was significantly increased ( $p = 0.007$ ) in the TGF- $\alpha$  + gastrin treated animals as compared to control animals; there was an approximately three-fold increase in pancreatic insulin content as compared to control animals. These data support the hypothesis that the combination of TFG- $\alpha$  and gastrin does produce an increase in the functional islet  $\beta$ -cell volume. This increase reflects an overall condition of  $\beta$ -cell hyperplasia (increase in number) rather than  $\beta$ -cell hypertrophy (increase in size of individual  $\beta$ -cells).

### EXAMPLE 5

#### Effect of Combination of TGF- $\alpha$ and Gastrin on Pancreatic

#### Insulin Content in Diabetic Animals

The second experiment was designed to determine whether the combination of TGF- $\alpha$  and gastrin could increase pancreatic insulin content in diabetic animals (streptozotocin (STZ) treated) to levels comparable to those in normal (non-STZ treated) animals.

Normal Wistar rats received a single iv injection of STZ at a dose of 80 mg/Kg body weight. This dose of STZ was intended to ensure that the study animals were rendered diabetic but that they retained a functioning but reduced  $\beta$ -cell mass. The STZ was dissolved immediately before administration in ice-cold 10 mM citric acid buffer. The animals were monitored daily; persistent diabetes was indicated by glycosuria and confirmed by non-fasting blood glucose determinations. One week after induction of diabetes, rats were randomly allocated into two groups (n = 6) as follows.

Group I: TGF- $\alpha$  + Gastrin: STZ diabetic rats were treated with a single i.p. injection of a combination of recombinant human TGF- $\alpha$  and synthetic rat Gastrin 1; both preparations were administered at a dose of 0.8  $\mu$ g/day for 10 days.

Group II: Control: STZ diabetic rats received an i.p. injection of vehicle alone for 10 days.

At the end of the study period, all animals were sacrificed and samples of pancreas taken and analyzed as described in Example 4 and the results are given in Table 2.

**Table 2**  
**Treatment of Streptozotocin Rats with TGF- $\alpha$  and Gastrin**

<u>Treatment</u>	<u>Pancreatic Insulin Content</u> ( $\mu$ g Insulin/mg protein)
Control (STZ alone)	6.06 $\pm$ 2.1
STZ plus TGF- $\alpha$ + Gastrin	26.7 $\pm$ 8.9

The induction of diabetes by STZ was successful and produced a moderate but

sustained degree of hyperglycemia. Total insulinopaenia was not sought so as to ensure that the study animals retained a functioning, but reduced  $\beta$ -cell mass.

As shown in Table 2, above, the pancreatic insulin content of the control streptozotocin treated animals was less than one third that of normal rats ( $20.6 \pm 6.0$  mg insulin/mg protein, *see* Table 1 above) as a result of destruction of  $\beta$ -cells by the STZ. In STZ animals treated with a combination of TGF- $\alpha$  and gastrin, the pancreatic insulin content was more than four-fold that of the animals which received STZ alone, and statistically the same as that of normal rats.

Diabetes mellitus is a disease in which the underlying physiological defect is a deficiency of  $\beta$ -cells as a result either of destruction of the  $\beta$ -cells due to auto-immune processes or of exhaustion of the potential for the  $\beta$ -cells to divide due to chronic stimulation from high circulating levels of glucose. The latter eventually leads to a situation when the process of  $\beta$ -cell renewal and/or replacement is compromised to the extent that there is an overall loss of  $\beta$ -cells and a concomitant decrease in the insulin content of the pancreas. The above results demonstrate that a combination of TGF- $\alpha$  and gastrin can be used to treat diabetes by stimulating the production of mature  $\beta$ -cells to restore the insulin content of the pancreas to non-diabetic levels.

#### EXAMPLE 6

##### Effects of TGF- $\alpha$ and Gastrin on IPGTT in STZ-Induced Diabetic Animals

Two groups (average body weight 103g) of STZ induced diabetic Wistar rats ( $n = 6/\text{group}$ ) were treated for 10 days with a daily i.p. injection of either a combination of TGF- $\alpha$  and gastrin or PBS. Fasting blood glucose was determined for all rats on days 0, 6, and 10. In order to establish that this insulin was secreted and functional, IPGTT tests were performed. At day 10, intraperitoneal glucose tolerance tests (IPGTT) were performed following an overnight fast. Blood samples were obtained from the tail vein, before and 30, 60 and 120 minutes after administration of an i.p. glucose injection at a dose of 2 g/kg body weight. Blood glucose determinations were performed as above. The blood glucose levels were similar in both study groups at time 0 but the TFG $\alpha$  and gastrin treated rats demonstrated a 50% reduction in blood glucose values (*see* Fig. 5), as compared to control rats at 30, 60, and 120 min. following the i.p. glucose load.

### EXAMPLE 7

#### Effects of TGF- $\alpha$ and Gastrin on Body Weight Gain and Insulin Content in Diabetes Prone Animals

5 Zucker rats were obtained at 30 days of age approximately 10-15 days prior to development of obesity. Besides the diabetes prone Zucker rats (genotype *fa/fa*, autosomal recessive mutation for obesity and diabetes), lean non-diabetic littermates (genotype *+/+*) also were included in the study as described below. The rats were monitored daily for development of obesity and diabetes by determining body weight and blood glucose. The onset of diabetes in Zucker rats usually started between days 45-50 and was confirmed by a significant increase in blood glucose levels, as compared to the levels in age-matched lean controls.

10 The study included 5 groups of 5 rats each as described in Table 3. Groups 1 and 2 (lean, non-diabetic) were treated with a TGF- $\alpha$  and gastrin combination or PBS respectively from day 0 to day 10. Groups 3, 4 and 5 included obese, early diabetic Zucker rats, genotype *fa/fa*. Group 3 received a combination pretreatment for 15 days (day -15 to day 0) prior to onset of diabetes and continuing post onset of diabetes for 10 additional days (day 0 to day 10). Group 4 was treated with a combination of TGF- $\alpha$  and gastrin for 10 days after onset of diabetes and Group 5 was treated with PBS over the same time period. At the end of the study, the rats were sacrificed and the pancreas removed. Small biopsies were taken from separate representative sites for protein and insulin determinations as described above.

20 The body weight gain in obese diabetic Zucker rats with pretreatment, treatment only or with saline (groups 3, 4, and 5 in Table 3) did not show any significant differences among the groups. It is interesting to note that even prolonged treatment (25 days, group 3) with TGF- $\alpha$  + gastrin was without effect on normal weight gain. Within error limits body weight gain was identical in all the groups.

25 The effect of TGF- $\alpha$  + gastrin treatment on fasting blood glucose in the obese Zucker rats was compared to the corresponding PBS controls. Fasting blood glucose was first significantly increased by day 15 ( $4.0 \pm 0.6$  vs.  $5.0 \pm 0.2$ ) and this time point was chosen as the starting time for the 10-day treatment period with TGF- $\alpha$  + gastrin or with PBS control. Fasting blood glucose levels were not significantly altered by the TGF- $\alpha$  + gastrin treatment or by PBS. Fasting blood glucose values were lower in lean, as compared to obese animals whether or not they were treated with the growth factors or with PBS.

Group	Geotype	Condition	<u>Pretreatment</u> + <u>Treatment (days)</u>	<u>PBS Control</u>	<u>Body Wt</u> <u>Gain (%±SE)</u>
1.	+/+	lean, non-diabetic	None	Yes	117±2.1
2.	+/+	lean, non-diabetic	0+10	No	119±1.9
3.	fa/fa	obese, early diabetic	-15+10	No	202±15
4.	fa/fa	obese, early diabetic	0+10	No	119±1.0
5.	fa/fa	obese, early diabetic	None	Yes	129±1.3

Group	Geotype	Condition	<u>Pretreatment</u> + <u>Treatment (days)</u>	<u>PBS Control</u>	<u>Body Wt</u> <u>Gain (%±SE)</u>
1.	+/+	lean, non-diabetic	None	Yes	117±2.1
2.	+/+	lean, non-diabetic	0+10	No	119±1.9
3.	fa/fa	obese, early diabetic	-15+10	No	202±15
4.	fa/fa	obese, early diabetic	0+10	No	119±1.0
5.	fa/fa	obese, early diabetic	None	Yes	129±1.3

1 The results of treatment with TGF- $\alpha$  and gastrin in the Zucker rat model of Type 2  
diabetes showed no significant differences in blood glucose levels between the treatment  
and control groups, probably reflecting the transient hypoglycemic effect following a  
prolonged period (18 hrs) of fasting. The immunohistochemical studies revealed significant  
5 increases in the number of single foci of insulin containing cells in the TGF- $\alpha$  and gastrin  
treated animals, as compared to control animals. These findings demonstrated an increase  
in single  $\beta$ -cells in adult rat pancreas following treatment with TGF- $\alpha$  and gastrin.  
Interestingly, such single  $\beta$ -cell foci are not commonly seen in adult (unstimulated) rat  
pancreas. These findings support a therapeutic role for TGF- $\alpha$  and gastrin in Type 1 and  
Type 2 diabetes since treatment is targeted at both  $\beta$ -cell neogenesis and replication.

10 The present invention is based in part on studies which demonstrated numerous  
insulin staining cells in the TGF- $\alpha$ -induced metaplastic ductules. The low level of exocrine  
and endocrine gene expression in the metaplastic ductal cells resembled that of  
15 protodifferentiated ductal cells seen in the early stage of fetal pancreatic development.  
Formation of islets (neogenesis) results from proliferation and differentiation of these  
protodifferentiated insulin expressing cells. Histologically this is manifested as islets  
appearing to bud from the pancreatic ducts (nesidioblastosis). In the MT-42 TGF- $\alpha$   
transgenic mice, ductular metaplasia was not seen in the immediate post-natal period, but  
20 only at 4 weeks of age. This indicates that TGF- $\alpha$  over-expression induced insulin  
expression in duct epithelia rather than prolonging the persistence of islet precursors found in  
fetal pancreatic ducts. Although the metaplastic ductules contained numerous insulin positive  
cells, the islet mass of the TGF- $\alpha$  transgenic mice was not increased over controls. The  
studies reported above demonstrate that complete islet cell neogenesis is reactivated *in vivo* in  
25 mammals in the ductular epithelium of the adult pancreas by stimulation with a gastrin/CCK  
receptor ligand, such as gastrin, and/or an EGF receptor ligand, such as TGF- $\alpha$ . Studies are  
reported on the transgenic over-expression of TGF- $\alpha$  and gastrin in the pancreas which  
elucidate the role of pancreatic gastrin expression in islet development and indicate that TGF-  
 $\alpha$  and gastrin each play a role in regulating islet development. Thus, regenerative  
30 differentiation of residual pluripotent pancreatic ductal cells into mature insulin-secreting  
cells is a viable method for the treatment of diabetes mellitus, by therapeutic administration

of this combination of factors or compositions which provide for their *in situ* expression within the pancreas.

**The present invention is not limited by the specific embodiments described herein.**

5        Modifications that become apparent from the foregoing description and accompanying figures fall within the scope of the claims.

Various publications are cited herein, the disclosures of which are incorporated by reference in their entirety.

[illegible]